

(1) DETERMINE THE STREAM-REACH BOUNDARY. (2) NEAR THE LOWER END OF THE REACH (IN THE DEEPEST PORTION OF THE RUN), COLLECT WATER SAMPLES AND ANALYZE USING THE CHEMICAL TESTS YOU HAVE AVAILABLE. YOU MAY USE YOUR COLLECTION CONTAINER TO OBSERVE WATERCOLOR AND CLARITY AND TO DETERMINE WATER ODORS. (3) MEASURE THE WIDTH-DEPTH AND VELOCITY, AND ESTIMATE THE WATER LEVEL. (4) IF YOU USE A TWO-POLE KICK-NET, COLLECT A MINIMUM OF THREE BENTHIC MACRO-INVERTEBRATE SAMPLES FROM THE BEST RIFFLE OR RUNS WITHIN YOUR STREAM REACH. USE THE TABLE ON PAGE FIVE TO RECORD INFORMATION ABOUT YOUR COLLECTIONS. (5) EVALUATE THE PHYSICAL AND HABITAT CONDITIONS; RECORD INFORMATION ABOUT KNOWN LAND USE ACTIVITIES. (6) SKETCH YOUR REACH OR SUBMIT PHOTOGRAPHS WITH THE SURVEY, AND ADD ANY OTHER COMMENTS THAT YOU FEEL ARE IMPORTANT. NOTE: A SCIENTIFIC COLLECTION PERMIT FROM THE WVDNR IS REQUIRED FOR ALL BENTHIC COLLECTIONS.

Stream name				ırvey date	
Watershed			Station	code	
Latitude	Longitude	Directi	ons to site		
Survey completed by					
Current weather cond					
Past weather condition	• • • • • • • • • • • • • • • • • • • •				
Affiliation		E-mail			
			Phone n	umber	
WATER CHEMISTRY: Usheets if necessary.	se the spaces below to record	the results of your wat	er chemistry ana	lysis; attach addi	ional
	Result units	Result	units	Res	ult units
Temperature (C/F)	Conduct		Alka		
Dissolved oxygen	Nitrate	es	Iro		
рН	Turbid	ity	Fecal	E-coli	
Additional tests (desc	ribe and record results)				
stream. The extra line condition; if so, be su	: Use the check boxes below to es are provided to write in any re to indicate these on your su ost dominant condition. If the	additional comments. `rvey (check all that ap	You may see mo ply). If multiple co	re than one type onditions are obse	of erved,
Water clarity	Water color	Water/Sedim	ent odor	Surface foam	
Clear	None	None		None	
Murky	Brown	Fishy		Slight	
Milky	Black	Musky		Moderate	
Muddy	Orange/red	Rotten egg		High	
Other (describe)	Gray/White	Sewage			
	Green	Chemical			
Algae color	Algae abundance	Algae growth	habit	Streambed color	
Light green	None	Even coati	ng	Brown	
Dark green	Scattered	Hairy		Black	
Brown	Moderate	Matted		Green	
Other (describe)	Heavy	Floating		White/gray	
				Orange/red	_
Physical condition co	mments:				

Estimate the percentage of your reach that is shaded.

> 80	80-60	60-40	< 40
Excellent	Good	Marginal	Poor

CIRCLE YOUR ESTIMATE

WIDTH AND DEPTH: Record the wetted width and depth of the channel's habitats (riffles, runs or pools). Choose two or more channel features to measure. Record the average depth from a minimum of four measurements (one of these should be from the deepest part of the habitat). The width should be measured from the widest section of the feature.

Riffle	Width (feet)	Depth (feet)
Run	Width (feet)	Depth (feet)
Pool	Width (feet)	Depth (feet)


CHANNEL PROFILES: Width and depth measurements can be used to create a cross section profile within your reach. Choose a location in your reach across one of the channel types above. Stretch a tape from bank to bank and anchor it at both ends. Move from left to right facing in an upstream direction; measure the distance from the stream bottom to the top of the tape at selected intervals (i.e. every foot). Record your measurements in the table below. The table provides enough spaces for 20 measurements; if more are necessary you can create your own table on a separate piece of paper. Your tape measure will probably not start at zero so make sure to record the actual position of the tape as you measure across the channel.

Width intervals

widin inter	vais								
1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
''	12	10	14	15	10	17	10	13	20
Depth mea	curamante								
Deputition	Suicilients		_	_		_			
1	2	3	4	5	6	1	8	9	10
11	12	13	14	15	16	17	18	19	20

PEBBLE COUNT: Collect a minimum of 100-particles from your reach using a Zigzag method, percent habitat method or specific transects (e.g. every 10-meter). If you do not complete a pebble count, **ALWAYS ESTIMATE** streambed composition from the riffles/runs chosen for your macroinvertebrate sample collections.

		Si	ze Classes (I	ntermediate a	xis in millimet	ers)	
Indicate your method from the choices below.	Silt/clay < 0.06	Sand 0.06 – 2	Fine Gravel 2 – 24	Coarse Gravel 25 – 64	Cobble 65 – 255	Boulder 256 – 1096	Bedrock > 1096
Zigzag							
% Habitat							
10-m Transects							
Woody Debris							
Includes sticks, roots,							
leaves etc.							
Totals							



- (A) Long axis (Length)
- (B) Intermediate axis (Width)
- (C) Short axis (Height)

Pebble counts require two people, one in the stream and one on shore. The person in the stream slowly walks upstream from bank to bank using one of the methods above. After each step the person reaches down without looking, picks up the first particle touched, and measures the intermediate axis with a ruler. The on-shore partner records the measurement. The process continues until 100 pebbles have been measured or the reach has been walked.

HABITAT CONDITIONS: Score each habitat condition using the scales provided. Add all of the scores to determine your overall habitat score and integrity rating. Feel free to describe additional features that you feel are important.

Sediment deposition	depo	ositio	o forn nal fea e rea	atures		dep	ositio 10% c	rease nal fea of the	atures	,	dep	ositio	nal fe	unts c atures reach	3;	dep	ositio	nounts n; > 6 affec	0% of	f
Score	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1

Embeddedness should be evaluated prior to or during your macroinvertebrate collections.

Embeddedness should be	e evaluated prior to or during	g your macroinvertebrate co	ollections.		
Embeddedness	Fine sediments surrounds <10% of the spaces between the gravel, cobble and	Fine sediment surrounds 10-30% of the spaces between the gravel, cobble and		Fine sediment surrounds > 60% of the spaces between the gravel, cobble and	
	boulders.	boulders.	boulders.	boulders.	
Score	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1	
Riffle frequency	Occurrence of riffles very frequent, making up > 60% of the reach; habitat variety is key in streams with continuous riffles; note the presence and/or absence of bends and other structures.	Occurrence of riffles relatively frequent, making up 40-60% of the reach; bends and/or other structures may provide additional habitat.	Occurrence of riffles infrequent, making up 20-40% of the reach; there are occasional riffles and bends, but the distances between such areas have greatly increased.	Occurrence of riffles < 20%; mostly all flat water throughout the reach; if riffles are present they are generally shallow and have very little cobble.	
Score	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6 5	4 3 2 1	
Attachment sites for invertebrates	Well developed riffles and runs; riffles are as wide as the stream and their lengths is twice that of the stream's width; cobble is prevalent, boulders and gravel also may be common.	Riffles are as wide as the stream but their length is less than twice the stream's width; cobble less abundant, gravel, boulders and other substrates maybe more common.	Riffle and runs lacking; if riffles are present they are not as wide as the stream nor is their length twice the stream's width; cobble rare, other substrate (esp. fines) more common.	Riffles and runs virtually nonexistent; large boulders and/or bedrock may be prevalent; or the reach may be mainly flat water throughout with finer sediments.	
Score	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6 5	4 3 2 1	
Velocity/depth regimes	All four velocity/depth combinations present; slow-shallow, slow - deep, fast-shallow and fast-deep.	3 of 4 velocity/depth combinations present; fast currents generally dominate (score lower if they are absent).	2 of 4 velocity/depth combinations present (score lower if fast current areas are absent).	Stream reach dominated by one velocity/depth regime (usually slow-shallow or slow-deep).	
Channel flow status	Water reaches the base of both lower banks and a minimal amount of channel substrate is exposed.	Water fills > 75% of the channel; < 25% of the channel substrate is exposed.	Water fills 25-75% of the channel; much of the riffle areas are exposed.	Very little water in the channel; mostly present as only standing pools.	
Score	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6 5	4 3 2 1	
Channel alterations	Stream straightening, dredging, artificial embankments, dams, bridge abutments etc. absent or minimal; stream has a natural pattern.	Some type of channel alteration present, usually in areas of bridge abutments; no evidence of recent channel alterations.	Artificial embankment structures present, at least to some extent, on both sides of the stream; 40-80% of the reach has been altered.	Stream banks are shored with gabions, rip/rap, cement etc.; > 80% of the reach has been disrupted.	
Score	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6 5	5 4 3 2 1	

The last three conditions are assessed on both sides of the channel. The **LEFT** and **RIGHT** sides are determined by looking downstream.

Bank ve prote	_	covered to vegetatio (trees, sh herbs) re- disruption	n; all levels rubs and presented; n from graz etc. minima Il plants	ing,	cove veg plar or n som veg	90% of the ered by natestation; one of the may be of well reported the disruption etation evices of the poton the ight reserved.	rural e level of missing resented; n of dent; > ential	covered to vegetation bare soil present a cropped v common;	n; patches	of contraction of the property	50% of the bovered by navegetation; dishiple, vegetationeen removed botential plantare greatly received.	tural sruption is on has I or the heights
Left		10	9	8	3	7	6	5	4	3	2	1
Right		10	9	8	3	7	6	5	4	3	2	1
Bank s	tability	evidence		or	stat of e	ks are mod ble; infreque rosion occu wn by bank r.	ent areas ur, mostly	unstable; reach has of erosior potential	e moderate 60% of the s some are n; high for erosion oding even	ely n e as as s b cats. 6	Banks are uns nany have ero areas (bare so straight section bends; obviou collapse or fail 50% of the rea erosion scars.	oded oils) along ns or s bank lure; >
Left		10	9	8	3	7	6	5	4	3	2	1
Right		10	9	8	3	7	6	5	4	3	2	1
Riparian bu	uffer width	vegetatio evidence impacts s parking lo clear-cuts	ots, road be	eds,	veg	e of undist etation 40- ne areas of urbance ev	60 ft;	vegetatio disturbed	indisturbed n 20-40 ft; areas throughout	the c	Zone of undist regetation < 2 disturbed area common throu entire reach.	0 ft; is
Left		10	9	8	3	7	6	5	4	3	2	1
Right		10	9	3	3	7	6	5	4	3	2	1

Total	Excellent	Very good	Good	Marginal	Poor
Score	> 170	170 - 150	149 - 130	129 - 100	< 100

SEDIMENT DEPOSITION MAY CAUSE THE FORMATION OF ISLANDS, POINT BARS (AREAS OF INCREASED DEPOSITION USUALLY AT THE BEGINNING OF A MEANDER THAT INCREASE IN SIZE AS THE CHANNEL IS DIVERTED TOWARD THE OUTER BANK) OR SHOALS, OR RESULT IN THE FILLING OF RUNS AND POOLS. USUALLY DEPOSITION IS EVIDENT IN AREAS THAT ARE OBSTRUCTED BY NATURAL OR MANMADE DEBRIS AND AREAS WHERE THE STREAM FLOW DECREASES, SUCH AS BENDS.

Habitat comments:			

LAND USE: Indicate the land uses that you believe may be having an impact on your stream station. Use the letters (**S**) streamside, (**M**) within ½ mile and (**W**) somewhere in the watershed, to indicate the approximate location of the disturbance and the numbers (**1**) slight, (**2**) moderate or (**3**) high, to represent the level of disturbance.

Active Construction	Pastureland	Single-family residences
Mountaintop mining	Cropland	Sub-urban developments
Deep mining	Intensive feedlots	Parking lots, strip-malls etc.
Abandoned mining	Unpaved Roads	Paved Roads
Logging	Trash dumps	Bridges
Oil and gas wells	Landfills	Other (describe)
Recreation (parks, trails etc.)	Industrial areas	

LEVEL-THREE SURVEY DATA SHEET Pipes? Yes Land use comments: No Describe the types of pipes observed and indicate if there is any discharge from the pipes. Also describe the color and odor of the discharge. **PHOTOGRAPH** and **SKETCH** YOUR **REACH**: Use the space below or a separate piece of paper to draw your study reach. Indicate the direction of flow, north, sample locations and important features of the reach. Photographs are an excellent method for tracking changes, especially changes related to the condition of the habitat. Choose a minimum of two permanent locations from which to take your photos. Submit your photos with your survey data sheet.

	BENTI	HIC COLL	ECTION DATA SHE			
Stream			Watershed			
Date	Station code		Latitude		Longitude	
Location				County		
		Г	7			
A 12 I .		Count	T . 1			Count
Annelida			Trichoptera			
	Total					
Bivalvia						
	Total					
Gastropoda						
					Tot	-1
			Plecoptera		Tot	ai
			ricooptora			
Constant	Total					
Crustacea						
	TOTAL				Tot	al
	101712		Odenete		Tot	ai
Ephemeroptera	1017.12		Odonata		100	aı
Ephemeroptera	101/12		Odonata		100	ai
Ephemeroptera	1017/12		Odonata		101	al
Ephemeroptera	101/10		Odonata		101	ai
Ephemeroptera	101/16		Odonata		100	al
Ephemeroptera	1017/12		Odonata			
Ephemeroptera			Odonata Coleoptera		Tot	
Ephemeroptera						
Ephemeroptera						
Ephemeroptera						
Ephemeroptera						
Ephemeroptera						
	Total					al
Ephemeroptera					Tot	al
			Coleoptera		Tot	al
	Total		Coleoptera		Tot	al
	Total		Coleoptera		Tot	al
Megaloptera	Total		Coleoptera		Tot	al
Megaloptera	Total		Coleoptera		Tot	al
Megaloptera	Total		Coleoptera		Tot	al
Megaloptera	Total		Coleoptera		Tot	al
Megaloptera	Total		Coleoptera		Tot	al
Megaloptera	Total		Coleoptera		Tot	al
Megaloptera	Total		Coleoptera		Tot	al

DISCHARGE: Determine the discharge by using a flow meter (if available) or other methods such as the FLOAT METHOD or the VELOCITY HEAD ROD METHOD (VHR). Discharge is measured from a run (area of the channel with fast moving water with no breaks in the surface such as protruding rocks). Indicate the methods chosen to measure the discharge and use the tables to record your results. Use the table to record your measurements.

Discharge method used			Water Level			
					1	
Float	L	Flow meter	Low	Normal	 High	Dry
Channel width		eet	LOW	Nomai	riigii	Diy
Charine width	'	CCI				
Distance (ft)	Depth (ft)	Velocity (ft/sec)	VHR (Rise-inches) Float (s		(sec) D	ischarge (cfs)
1	, , ,	• , , ,	,		,	<u> </u>
2						
3						
4						
5						
6						
7						
8						
9						
10						
11						
12						
13						
14						
15						
16						
17						
18						
19						
20						
	1		1	'	•	
Average Dep	oth	feet		pelow to determin		
				d above. The ris		
Cross Sectional Area (CSA) ft ²			Rise (R)	Velocity	Rise (R)	Velocity
(CSA = Average Depth x V	Vidth)		1/4	1.2	3 1/4	4.2
			1/2	1.6	3 ½	4.3
			3/4	2.0	3 3/4	4.5
Discharge = CSA x Velocity			1 1/4	2.3 2.6	4 1/4	4.6 4.8
- · · · · · · · · · · · · · · · · · · ·			1 1/2	2.8	4 1/2	4.8
= x = cfs (ft ³ /sec)			1 3/4	3.1	4 1/2	5.0
	2	3.3	5	5.2		
If you use a float	2 1/4	3.5	5 1/4	5.3		
number of seconds it took to travel the distance in the			2 1/2	3.7	5 ½	5.4
column indicated.			2 3/4	3.8	5 3/4	5.5
			3	4.0	6	5.7

Submit the <u>original or a clear copy</u> of your survey to the **Citizen's Monitoring Coordinator** at the address below. The information will be reviewed and returned.

WV DEPARTMENT OF ENVIRONMENTAL PROTECTION
SAVE OUR STREAMS PROGRAM
601 57TH STREET, SE
CHARLESTON, WV 25304

For more information go to: http://www.dep.wv.gov/sos

Float distance (feet) _____